

Exploiting Structured Sparsity in Large Scale Semidefinite Programming Problems

M. Kojima
Tokyo Institute of Technology

November 2010

At National Cheng-Kung University, Tainan

- Kim, Kojima, Mevissen and Yamashita, “Exploiting sparsity in linear and nonlinear inequalities via positive semidefinite matrix completion”, *Mathematical Programming* to appear.

.. - p.1/35

Outline

- 0 Semidefinite Programming (SDP)
- 1 A simple example for 2 types of sparsities
- 2 Chordal graph
- 3 Domain-space sparsity
- 4 Range-space sparsity
- 5 Numerical results
- 6 Concluding remarks

.. - p.2/35

Outline

0 Semidefinite Programming (SDP)

- 1 A simple example for 2 types of sparsities
- 2 Chordal graph
- 3 Domain-space sparsity
- 4 Range-space sparsity
- 5 Numerical results
- 6 Concluding remarks

.. - p.3/35

A general linear (or nonlinear) SDP

= “Optimization problem involving an $n \times n$ real symmetric matrix variable X to be positive semidefinite”

.. - p.4/35

A general linear (or nonlinear) SDP

= “Optimization problem involving an $n \times n$ real symmetric matrix variable X to be positive semidefinite”

min. a linear (or nonlinear) function in $y \in \mathbb{R}^m$, $X \in \mathbb{S}^n$,
 sub. to linear (or nonlinear) equalities and inequalities
 in $y \in \mathbb{R}^m$, $X \in \mathbb{S}^n$,

$$X = \begin{pmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \dots & \dots & \dots & \dots \\ X_{n1} & X_{n2} & \dots & X_{nn} \end{pmatrix} \succeq O$$

(positive semidefinite).

Here \mathbb{S}^n denotes the space of $n \times n$ symmetric matrices.

- We can solve linear SDP by interior-point methods.
- We will discuss 2 types of conversions of a large-scale SDP satisfying a structured sparsity to solve it efficiently.

.. p.4/35

Applications of SDPs

- System and control theory — Linear matrix inequality
- Robust Optimization
- Machine learning
- Quantum chemistry
- Quantum computation
- Moment problems (Applied probability)
- SDP relaxation —
 Max cut, Max clique, Sensor network localization,
 Polynomial optimization
- Design optimization of structures
- ...

In many applications, SDPs are large-scale and often satisfy a certain sparsity characterized by a chordal graph structure.

.. p.5/35

Outline

- 0 Semidefinite Programming (SDP)
- 1 A simple example for 2 types of sparsities
- 2 Chordal graph
- 3 Domain-space sparsity
- 4 Range-space sparsity
- 5 Numerical results
- 6 Concluding remarks

.. p.6/35

$$\text{SDP: } \min \sum_{i=1}^{n-1} (X_{ii} + b_i(X_{i,i+1} + X_{i+1,i})) + X_{nn} \quad (1)$$

sub. to (Matrix inequality, diagonal+bordered)

$$M(X) = \begin{pmatrix} 1 - X_{11} & 0 & \dots & X_{12} \\ 0 & 1 - X_{22} & \dots & X_{23} \\ \dots & \dots & \ddots & \dots \\ X_{21} & X_{32} & \dots & 1 - X_{nn} \end{pmatrix} \succeq O \quad (2)$$

$$X = \begin{pmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \dots & \dots & \dots & \dots \\ X_{n1} & X_{n2} & \dots & X_{nn} \end{pmatrix} \succeq O \quad (\text{positive semidefinite})$$

- The number of variables is $n(n+1)/2$; $X_{ij} = X_{ji}$.
- domain-space sparsity — Only X_{ij} ($|i-j| \leq 1$) are used in (1), (2) among all variables X_{ij} ($1 \leq i \leq j \leq n$).
- range-space sparsity — (2) is diagonal + bordered.

.. p.7/35

$$\text{SDP: } \min \sum_{i=1}^{n-1} (X_{ii} + b_i(X_{i,i+1} + X_{i+1,i})) + X_{nn} \quad (1)$$

sub. to (Matrix inequality, diagonal+bordered)

$$M(\mathbf{X}) = \begin{pmatrix} 1 - X_{11} & 0 & \dots & X_{12} \\ 0 & 1 - X_{22} & \dots & X_{23} \\ \dots & \dots & \ddots & \dots \\ X_{21} & X_{32} & \dots & 1 - X_{nn} \end{pmatrix} \succeq \mathbf{O} \quad (2)$$

$$\mathbf{X} = \begin{pmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \dots & \dots & \dots & \dots \\ X_{n1} & X_{n2} & \dots & X_{nn} \end{pmatrix} \succeq \mathbf{O} \quad (\text{positive semidefinite})$$

.. p.8/35

$$\text{SDP: } \min \sum_{i=1}^{n-1} (X_{ii} + b_i(X_{i,i+1} + X_{i+1,i})) + X_{nn} \quad (1)$$

sub. to (Matrix inequality, diagonal+bordered)

$$M(\mathbf{X}) = \begin{pmatrix} 1 - X_{11} & 0 & \dots & X_{12} \\ 0 & 1 - X_{22} & \dots & X_{23} \\ \dots & \dots & \ddots & \dots \\ X_{21} & X_{32} & \dots & 1 - X_{nn} \end{pmatrix} \succeq \mathbf{O} \quad (2)$$

$$\mathbf{X} = \begin{pmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \dots & \dots & \dots & \dots \\ X_{n1} & X_{n2} & \dots & X_{nn} \end{pmatrix} \succeq \mathbf{O} \quad (\text{positive semidefinite})$$

↓ conversion with exploiting the domain and range sparsities
 “smaller size” SDP equivalent to the original SDP

- Next, numerical results on the converted SDP
- Later, technical details on the conversion = the subject of this talk

.. p.8/35

Numerical results

- SeDuMi (MATLAB, a prima-dual interior-point method)
- 2.66 GHz Dual-Core Intel Xeon with 12GB memory

size of \mathbf{X} = n	SeDuMi elapsed time (second)	
	Original SDP	Converted SDP with exploiting d-space & r-space sparsities
10	0.2	0.1
100	1091.4	0.6
1000	-	6.3
10000	-	99.2

.. p.9/35

Outline



- 0 Semidefinite Programming (SDP)
- 1 A simple example for 2 types of sparsities
- 2 Chordal graph
- 3 Domain-space sparsity
- 4 Range-space sparsity
- 5 Numerical results
- 6 Concluding remarks

- Sparsity pattern will be described in terms of a graph.
- We will assume that the sparsity pattern graph has a sparse chordal extension to exploit the domain- and range-space sparsity in SDPs.

.. p.10/35


$G(N, E)$: a graph, $N = \{1, \dots, n\}$ (nodes), $E \subset N \times N$ (edges)

chordal $\Leftrightarrow \forall$ cycle with more than 3 edges has a chord




Sparsity pattern is described in terms of a graph

$$R = \begin{pmatrix} * & 0 & 0 & 0 & 0 & * \\ 0 & * & 0 & 0 & 0 & * \\ 0 & 0 & * & * & 0 & * \\ 0 & 0 & * & * & * & 0 \\ 0 & 0 & 0 & * & * & * \\ * & * & * & 0 & * & * \end{pmatrix}$$



$$R = \begin{pmatrix} * & 0 & 0 & 0 & 0 & * \\ 0 & * & 0 & 0 & 0 & * \\ 0 & 0 & * & * & 0 & * \\ 0 & 0 & * & * & * & * \\ 0 & 0 & 0 & * & * & * \\ * & * & * & * & * & * \end{pmatrix}$$



Outline

- 0 Semidefinite Programming (SDP)
- 1 A simple example for 2 types of sparsities
- 2 Chordal graph
- 3 Domain-space sparsity
- 4 Range-space sparsity
- 5 Numerical results
- 6 Concluding remarks

Opt. problem involving a symmetric matrix variable $\mathbf{X} \succeq \mathbf{O}$:

$$(P) \min f_0(\mathbf{y}, \mathbf{X}) \text{ sub.to } \mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega, \mathbb{S}^n \ni \mathbf{X} \succeq \mathbf{O}.$$

Here $f_0 : \mathbb{R}^s \times \mathbb{S}^n \rightarrow \mathbb{R}$, $\mathbf{f} : \mathbb{R}^s \times \mathbb{S}^n \rightarrow V \supset \Omega$.

d-space sparsity pattern graph $G(N, F)$: $N = \{1, 2, \dots, n\}$,

$$F = \left\{ (i, j) : \begin{array}{l} i \neq j, X_{ij} \text{ is necessary} \\ \text{to evaluate } f_0(\mathbf{y}, \mathbf{X}) \text{ or } \mathbf{f}(\mathbf{y}, \mathbf{X}) \end{array} \right\}$$

Opt. problem involving a symmetric matrix variable $\mathbf{X} \succeq \mathbf{O}$:

(P) min $f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega$, $\mathbb{S}^n \ni \mathbf{X} \succeq \mathbf{O}$.

Here $f_0 : \mathbb{R}^s \times \mathbb{S}^n \rightarrow \mathbb{R}$, $\mathbf{f} : \mathbb{R}^s \times \mathbb{S}^n \rightarrow V \supset \Omega$.

d-space sparsity pattern graph $G(N, F)$: $N = \{1, 2, \dots, n\}$,

$$F = \left\{ (i, j) : \begin{array}{l} i \neq j, X_{ij} \text{ is necessary} \\ \text{to evaluate } f_0(\mathbf{y}, \mathbf{X}) \text{ or } \mathbf{f}(\mathbf{y}, \mathbf{X}) \end{array} \right\}$$

$$\min f_0(\mathbf{y}, \mathbf{X}) = \sum_{i=1}^3 (y_i X_{ii} + X_{i,i+1} + X_{i+1,i})$$

sub. to

$$\mathbf{f}(\mathbf{y}, \mathbf{X}) = \begin{pmatrix} 1 - X_{11} & X_{12} & y_1 & 2y_2 \\ X_{21} & 1 - X_{22} & X_{23} & 3y_3 \\ y_1 & X_{32} & 1 - X_{33} & X_{34} \\ 2y_2 & 3y_3 & X_{43} & 1 - X_{44} \end{pmatrix} \succeq \mathbf{O},$$

$$\mathbb{S}^4 \ni \mathbf{X} \succeq \mathbf{O}$$

...p.14/35

Opt. problem involving a symmetric matrix variable $\mathbf{X} \succeq \mathbf{O}$:

(P) min $f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega$, $\mathbb{S}^n \ni \mathbf{X} \succeq \mathbf{O}$.

Here $f_0 : \mathbb{R}^s \times \mathbb{S}^n \rightarrow \mathbb{R}$, $\mathbf{f} : \mathbb{R}^s \times \mathbb{S}^n \rightarrow V \supset \Omega$.

d-space sparsity pattern graph $G(N, F)$: $N = \{1, 2, \dots, n\}$,

$$F = \left\{ (i, j) : \begin{array}{l} i \neq j, X_{ij} \text{ is necessary} \\ \text{to evaluate } f_0(\mathbf{y}, \mathbf{X}) \text{ or } \mathbf{f}(\mathbf{y}, \mathbf{X}) \end{array} \right\}$$

$$\min f_0(\mathbf{y}, \mathbf{X}) = \sum_{i=1}^3 (y_i X_{ii} + X_{i,i+1} + X_{i+1,i})$$

sub. to

$$\mathbf{f}(\mathbf{y}, \mathbf{X}) = \begin{pmatrix} 1 - X_{11} & X_{12} & y_1 & 2y_2 \\ X_{21} & 1 - X_{22} & X_{23} & 3y_3 \\ y_1 & X_{32} & 1 - X_{33} & X_{34} \\ 2y_2 & 3y_3 & X_{43} & 1 - X_{44} \end{pmatrix} \succeq \mathbf{O},$$

$$\mathbb{S}^4 \ni \mathbf{X} \succeq \mathbf{O} \quad \Rightarrow N = \{1, 2, 3, 4\}$$

- X_{ij} , $|i - j| \leq 1$ are necessary to evaluate $f_0(\mathbf{y}, \mathbf{X})$, $\mathbf{f}(\mathbf{y}, \mathbf{X})$
- $F = \{(i, i + 1) : i = 1, 2, 3\}$

$G(N, F)$ = a chordal graph $\textcircled{1} - \textcircled{2} - \textcircled{3} - \textcircled{4}$

...p.14/35

Opt. problem involving a symmetric matrix variable $\mathbf{X} \succeq \mathbf{O}$:

(P) min $f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega$, $\mathbb{S}^n \ni \mathbf{X} \succeq \mathbf{O}$.

Here $f_0 : \mathbb{R}^s \times \mathbb{S}^n \rightarrow \mathbb{R}$, $\mathbf{f} : \mathbb{R}^s \times \mathbb{S}^n \rightarrow V \supset \Omega$.

d-space sparsity pattern graph $G(N, F)$: $N = \{1, 2, \dots, n\}$,

$$F = \left\{ (i, j) : \begin{array}{l} i \neq j, X_{ij} \text{ is necessary} \\ \text{to evaluate } f_0(\mathbf{y}, \mathbf{X}) \text{ or } \mathbf{f}(\mathbf{y}, \mathbf{X}) \end{array} \right\}$$

...p.15/35

Opt. problem involving a symmetric matrix variable $\mathbf{X} \succeq \mathbf{O}$:

(P) min $f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega$, $\mathbb{S}^n \ni \mathbf{X} \succeq \mathbf{O}$.

Here $f_0 : \mathbb{R}^s \times \mathbb{S}^n \rightarrow \mathbb{R}$, $\mathbf{f} : \mathbb{R}^s \times \mathbb{S}^n \rightarrow V \supset \Omega$.

d-space sparsity pattern graph $G(N, F)$: $N = \{1, 2, \dots, n\}$,

$$F = \left\{ (i, j) : \begin{array}{l} i \neq j, X_{ij} \text{ is necessary} \\ \text{to evaluate } f_0(\mathbf{y}, \mathbf{X}) \text{ or } \mathbf{f}(\mathbf{y}, \mathbf{X}) \end{array} \right\}$$

$$\Updownarrow \begin{array}{l} G(N, E) : \text{a chordal extension of } G(N, F) \\ C_1, C_2, \dots, C_\ell : \text{the maximal cliques of } G(N, E) \end{array}$$

(P') min $f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega$, $\mathbf{X}(C_p) \succeq \mathbf{O}$ ($p = 1, \dots, \ell$).

Here $\mathbf{X}(C_p)$: a submatrix consisting of X_{ij} , $(i, j) \in C_p \times C_p$.

...p.15/35

Opt. problem involving a symmetric matrix variable $\mathbf{X} \succeq \mathbf{O}$:

(P) $\min f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega, \mathbb{S}^n \ni \mathbf{X} \succeq \mathbf{O}$.

Here $f_0: \mathbb{R}^s \times \mathbb{S}^n \rightarrow \mathbb{R}, \mathbf{f}: \mathbb{R}^s \times \mathbb{S}^n \rightarrow V \supset \Omega$.


d-space sparsity pattern graph $G(N, F): N = \{1, 2, \dots, n\}$,

$$F = \left\{ (i, j) : \begin{array}{l} i \neq j, X_{ij} \text{ is necessary} \\ \text{to evaluate } f_0(\mathbf{y}, \mathbf{X}) \text{ or } \mathbf{f}(\mathbf{y}, \mathbf{X}) \end{array} \right\}$$

$$\Updownarrow \left\{ \begin{array}{l} G(N, E) : \text{a chordal extension of } G(N, F) \\ C_1, C_2, \dots, C_\ell : \text{the maximal cliques of } G(N, E) \end{array} \right.$$

(P') $\min f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega, \mathbf{X}(C_p) \succeq \mathbf{O} (p = 1, \dots, \ell)$.

Here $\mathbf{X}(C_p)$: a submatrix consisting of $X_{ij}, (i, j) \in C_p \times C_p$.



.. p.15/35

Opt. problem involving a symmetric matrix variable $\mathbf{X} \succeq \mathbf{O}$:

(P) $\min f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega, \mathbb{S}^n \ni \mathbf{X} \succeq \mathbf{O}$.

Here $f_0: \mathbb{R}^s \times \mathbb{S}^n \rightarrow \mathbb{R}, \mathbf{f}: \mathbb{R}^s \times \mathbb{S}^n \rightarrow V \supset \Omega$.

d-space sparsity pattern graph $G(N, F): N = \{1, 2, \dots, n\}$,

$$F = \left\{ (i, j) : \begin{array}{l} i \neq j, X_{ij} \text{ is necessary} \\ \text{to evaluate } f_0(\mathbf{y}, \mathbf{X}) \text{ or } \mathbf{f}(\mathbf{y}, \mathbf{X}) \end{array} \right\}$$

$$\Updownarrow \left\{ \begin{array}{l} G(N, E) : \text{a chordal extension of } G(N, F) \\ C_1, C_2, \dots, C_\ell : \text{the maximal cliques of } G(N, E) \end{array} \right.$$

(P') $\min f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega, \mathbf{X}(C_p) \succeq \mathbf{O} (p = 1, \dots, \ell)$.

Here $\mathbf{X}(C_p)$: a submatrix consisting of $X_{ij}, (i, j) \in C_p \times C_p$.

.. p.16/35

Opt. problem involving a symmetric matrix variable $\mathbf{X} \succeq \mathbf{O}$:

(P) $\min f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega, \mathbb{S}^n \ni \mathbf{X} \succeq \mathbf{O}$.

Here $f_0: \mathbb{R}^s \times \mathbb{S}^n \rightarrow \mathbb{R}, \mathbf{f}: \mathbb{R}^s \times \mathbb{S}^n \rightarrow V \supset \Omega$.

d-space sparsity pattern graph $G(N, F): N = \{1, 2, \dots, n\}$,

$$F = \left\{ (i, j) : \begin{array}{l} i \neq j, X_{ij} \text{ is necessary} \\ \text{to evaluate } f_0(\mathbf{y}, \mathbf{X}) \text{ or } \mathbf{f}(\mathbf{y}, \mathbf{X}) \end{array} \right\}$$

$$\Updownarrow \left\{ \begin{array}{l} G(N, E) : \text{a chordal extension of } G(N, F) \\ C_1, C_2, \dots, C_\ell : \text{the maximal cliques of } G(N, E) \end{array} \right.$$

(P') $\min f_0(\mathbf{y}, \mathbf{X})$ sub.to $\mathbf{f}(\mathbf{y}, \mathbf{X}) \in \Omega, \mathbf{X}(C_p) \succeq \mathbf{O} (p = 1, \dots, \ell)$.

Here $\mathbf{X}(C_p)$: a submatrix consisting of $X_{ij}, (i, j) \in C_p \times C_p$.

- (P) \Leftrightarrow (P') is based on the positive definite matrix completion (Grone et al. 1984).

.. p.16/35

Outline

- 0 Semidefinite Programming (SDP)
- 1 A simple example for 2 types of sparsities
- 2 Chordal graph
- 3 Domain-space sparsity
- 4 Range-space sparsity
- 5 Numerical results
- 6 Concluding remarks

.. p.17/35

$G(N, E)$: a chordal graph with $N = \{1, \dots, n\}$ and the max. cliques of C_1, \dots, C_ℓ . $E^\bullet = E \cup \{(i, i) : i \in N\}$.
 $\mathbb{S}^n(E^\bullet) = \{\mathbf{Y} \in \mathbb{S}^n : Y_{ij} = 0 \text{ if } (i, j) \notin E^\bullet\}$.
 $\mathbb{S}_+^C = \{\mathbf{Y} \succeq \mathbf{O} : Y_{ij} = 0 \text{ if } (i, j) \notin C \times C\} \text{ for } \forall C \subseteq N$.

Theorem (Agler, Helton, McCulough and Rodman 1988)
 Suppose $\mathbf{M} \in \mathbb{S}^n(E^\bullet)$. $\mathbf{M} \succeq \mathbf{O}$ iff
 $\mathbf{M} = \mathbf{Y}^1 + \mathbf{Y}^2 + \dots + \mathbf{Y}^\ell$ for $\exists \mathbf{Y}^k \in \mathbb{S}_+^{C_k}$ ($k = 1, \dots, \ell$).

$G(N, E)$: a chordal graph with $N = \{1, \dots, n\}$ and the max. cliques of C_1, \dots, C_ℓ . $E^\bullet = E \cup \{(i, i) : i \in N\}$.
 $\mathbb{S}^n(E^\bullet) = \{\mathbf{Y} \in \mathbb{S}^n : Y_{ij} = 0 \text{ if } (i, j) \notin E^\bullet\}$.
 $\mathbb{S}_+^C = \{\mathbf{Y} \succeq \mathbf{O} : Y_{ij} = 0 \text{ if } (i, j) \notin C \times C\} \text{ for } \forall C \subseteq N$.

Theorem (Agler, Helton, McCulough and Rodman 1988)
 Suppose $\mathbf{M} \in \mathbb{S}^n(E^\bullet)$. $\mathbf{M} \succeq \mathbf{O}$ iff
 $\mathbf{M} = \mathbf{Y}^1 + \mathbf{Y}^2 + \dots + \mathbf{Y}^\ell$ for $\exists \mathbf{Y}^k \in \mathbb{S}_+^{C_k}$ ($k = 1, \dots, \ell$).

①—②—③ $C_1 = \{1, 2\}$, $C_2 = \{2, 3\}$. $\mathbf{M} : \mathbb{R}^m \rightarrow \mathbb{S}^3(E^\bullet)$.

$$\mathbf{M}(\mathbf{u}) = \begin{pmatrix} M_{11}(\mathbf{u}) & M_{12}(\mathbf{u}) & 0 \\ M_{21}(\mathbf{u}) & M_{22}(\mathbf{u}) & M_{23}(\mathbf{u}) \\ 0 & M_{32}(\mathbf{u}) & M_{33}(\mathbf{u}) \end{pmatrix}$$

$G(N, E)$: a chordal graph with $N = \{1, \dots, n\}$ and the max. cliques of C_1, \dots, C_ℓ . $E^\bullet = E \cup \{(i, i) : i \in N\}$.
 $\mathbb{S}^n(E^\bullet) = \{\mathbf{Y} \in \mathbb{S}^n : Y_{ij} = 0 \text{ if } (i, j) \notin E^\bullet\}$.
 $\mathbb{S}_+^C = \{\mathbf{Y} \succeq \mathbf{O} : Y_{ij} = 0 \text{ if } (i, j) \notin C \times C\} \text{ for } \forall C \subseteq N$.

Theorem (Agler, Helton, McCulough and Rodman 1988)
 Suppose $\mathbf{M} \in \mathbb{S}^n(E^\bullet)$. $\mathbf{M} \succeq \mathbf{O}$ iff
 $\mathbf{M} = \mathbf{Y}^1 + \mathbf{Y}^2 + \dots + \mathbf{Y}^\ell$ for $\exists \mathbf{Y}^k \in \mathbb{S}_+^{C_k}$ ($k = 1, \dots, \ell$).

①—②—③ $C_1 = \{1, 2\}$, $C_2 = \{2, 3\}$. $\mathbf{M} : \mathbb{R}^m \rightarrow \mathbb{S}^3(E^\bullet)$.

$\mathbf{M}(\mathbf{u}) \succeq \mathbf{O}$

$G(N, E)$: a chordal graph with $N = \{1, \dots, n\}$ and the max. cliques of C_1, \dots, C_ℓ . $E^\bullet = E \cup \{(i, i) : i \in N\}$.
 $\mathbb{S}^n(E^\bullet) = \{\mathbf{Y} \in \mathbb{S}^n : Y_{ij} = 0 \text{ if } (i, j) \notin E^\bullet\}$.
 $\mathbb{S}_+^C = \{\mathbf{Y} \succeq \mathbf{O} : Y_{ij} = 0 \text{ if } (i, j) \notin C \times C\} \text{ for } \forall C \subseteq N$.

Theorem (Agler, Helton, McCulough and Rodman 1988)
 Suppose $\mathbf{M} \in \mathbb{S}^n(E^\bullet)$. $\mathbf{M} \succeq \mathbf{O}$ iff
 $\mathbf{M} = \mathbf{Y}^1 + \mathbf{Y}^2 + \dots + \mathbf{Y}^\ell$ for $\exists \mathbf{Y}^k \in \mathbb{S}_+^{C_k}$ ($k = 1, \dots, \ell$).

①—②—③ $C_1 = \{1, 2\}$, $C_2 = \{2, 3\}$. $\mathbf{M} : \mathbb{R}^m \rightarrow \mathbb{S}^3(E^\bullet)$.

$$\mathbf{M}(\mathbf{u}) \succeq \mathbf{O} \iff \mathbf{M}(\mathbf{u}) = \begin{pmatrix} \mathbf{Y}_{11}^1 & \mathbf{Y}_{12}^1 & 0 \\ \mathbf{Y}_{12}^1 & \mathbf{Y}_{22}^1 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & \mathbf{Y}_{22}^2 & \mathbf{Y}_{23}^2 \\ 0 & \mathbf{Y}_{32}^2 & \mathbf{Y}_{33}^2 \end{pmatrix}$$

$$\iff \left\{ \begin{array}{l} M_{11} = \mathbf{Y}_{11}^1, M_{12} = \mathbf{Y}_{12}^1, \\ M_{22} = \mathbf{Y}_{22}^1 + \mathbf{Y}_{22}^2, \\ M_{23} = \mathbf{Y}_{23}^2, M_{33} = \mathbf{Y}_{33}^2, \\ \square \succeq \mathbf{O}, \square \succeq \mathbf{O} \end{array} \right\} \iff \left\{ \begin{array}{l} \begin{pmatrix} M_{11}(\mathbf{u}) & M_{12}(\mathbf{u}) \\ M_{21}(\mathbf{u}) & \mathbf{Y}_{22}^1 \end{pmatrix} \succeq \mathbf{O}, \\ \begin{pmatrix} M_{22}(\mathbf{u}) - \mathbf{Y}_{22}^1 & M_{23}(\mathbf{u}) \\ M_{32}(\mathbf{u}) & M_{33}(\mathbf{u}) \end{pmatrix} \succeq \mathbf{O} \end{array} \right.$$

Summary of the d-space and r-space conversion methods:

Sparsity characterized by a chordal graph structure

SDP (linear, polynomial, nonlinear)
each large-scale matrix variable
↓ exploiting d-space sparsity
multiple smaller matrix variables
each large-scale matrix inequality
↓ exploiting r-space sparsity
multiple smaller matrix inequalities

→ SparseCoLO
for linear SDP

↓ if SDP is linear ↓ relaxation if SDP is polynomial

Linear SDP with multiple smaller matrix variables and matrix inequalities

.. p.20/35

Summary of the d-space and r-space conversion methods:

Sparsity characterized by a chordal graph structure

SDP (linear, polynomial, nonlinear)
each large-scale matrix variable
↓ exploiting d-space sparsity
multiple smaller matrix variables
each large-scale matrix inequality
↓ exploiting r-space sparsity
multiple smaller matrix inequalities

→ SparseCoLO
for linear SDP

↓ if SDP is linear ↓ relaxation if SDP is polynomial

Linear SDP with multiple smaller matrix variables and matrix inequalities

● SparsePOP = sparse SDP relaxation (Waki et. al '06) :

POP ⇒ Poly. SDP ⇒ Linear SDP
adding valid poly. mat. inequalities ← sparsity relaxation

.. p.20/35

Outline

- 0 Semidefinite Programming (SDP)
- 1 A simple example for 2 types of sparsities
- 2 Chordal graph
- 3 Domain-space sparsity
- 4 Range-space sparsity
- 5 Numerical results
- 6 Concluding remarks

.. p.21/35

Test Problems

- (a) SDP relaxation of quadratic optimization problems (QOPs)
- (b) Linear SDP relaxation of randomly generated sparse quadratic SDPs
- (c) Polynomial optimization problems (POPs)

- We apply SparseCoLO+ SDPA to (a) and (b), where SparseCoLO — MATLAB software for the d-space and r-space conversion methods, SDPA — a primal-dual interior-point method for SDPs.
- We apply SparsePOP + SDPA to (c), where SparsePOP — a sparse SDP relaxation for POPs using the d-space conversion method.
- 3.06 GHz Intel Core 2 Duo with 8 GB memory.



.. p.22/35

(a) Linear SDP relaxation of **sparse QOPs**

Sparse Linear SDP	size X	No. of equalities	E. time in seconds	
			no sparsity	d-space
M1000.05	1000	1000	41.2	0.5
M1000.15	1000	1000	39.6	52.7
thetaG11	801	2401	41.8	6.9
qpG11	1600	800	112.5	3.1
sensor1000	1002	11010	271.8	18.3
sensor4000	4002	47010	o.mem.	56.0

Sparse Linear SDP **sparse QOP**
M1000.?? \Leftarrow max cut problems with diff. edge densities
thetaG11 \Leftarrow minimization of the Lovasz theta function
qpG11 \Leftarrow a box constrained QOP
sensor???? \Leftarrow a sensor network localization problem with ???? sensors

M1000.05





d-space sparsity pattern

d-space sparsity pattern with the symmetric min. deg. ordering (symamd, MATLAB)

Before conversion \Rightarrow After conversion
one $1000 \times 1000 X \succeq O$ 117 smaller $X_k \succeq O$
max. size = 31×31
ave. size = 10.1×10.1
41.5 second 0.5 second

M1000.15





d-space sparsity pattern

d-space sparsity pattern with the symmetric min. deg. ordering (symamd, MATLAB)

Before conversion \Rightarrow After conversion
one $1000 \times 1000 X \succeq O$ 47 smaller $X_k \succeq O$
max. size = 91×91
ave. size = 36.6×36.6
39.6 second 52.5 second

sensor1000





d-space sparsity pattern

d-space sparsity pattern with the symmetric min. deg. ordering (symamd, MATLAB)

Before conversion \Rightarrow After conversion
one $1002 \times 1002 X \succeq O$ 914 smaller $X_k \succeq O$
max. size = 34×34
ave. size = 6.2×6.2
271.3 second 18.3 second

sensor4000



d-space sparsity pattern

d-space sparsity pattern with the symmetric min. deg. ordering (symamd, MATLAB)

Before conversion \Rightarrow After conversion
 one 4002×4002 $\mathbf{X} \succeq \mathbf{O}$ 3892 smaller $\mathbf{X}_k \succeq \mathbf{O}$
 out of memory max. size = 37×37 ,
 ave. size = 5.3×5.3
 56.0

(b) Linear SDP relaxation of a sparse quadratic SDP

Quadratic SDP: $\min c^T x$ sub to $M(x) \succeq \mathbf{O}$,

where $M : \mathbb{R}^s \rightarrow \mathbb{S}^n$ whose (i, j) element is given by

$$M_{ij}(x) = (1, x^T) Q_{ij} \begin{pmatrix} 1 \\ x \end{pmatrix} = Q_{ij} \bullet \begin{pmatrix} 1 & x^T \\ x & xx^T \end{pmatrix}, \forall x \in \mathbb{R}^s.$$

Here $Q \bullet Y = \text{trace } Q^T Y$ (the inner product of Q and Y).

(b) Linear SDP relaxation of a sparse quadratic SDP

SDP: $\min c^T x$ sub to $\widehat{M}(x, \mathbf{X}) \succeq \mathbf{O}, \begin{pmatrix} x_0 & x^T \\ x & \mathbf{X} \end{pmatrix} \succeq \mathbf{O}, x_0 = 1$

where $\widehat{M} : \mathbb{R}^s \times \mathbb{S}^s \rightarrow \mathbb{S}^n$ whose (i, j) element is given by

$$\widehat{M}_{ij}(x, \mathbf{X}) = Q_{ij} \bullet \begin{pmatrix} 1 & x^T \\ x & \mathbf{X} \end{pmatrix} \text{ for every } x \in \mathbb{R}^s, \mathbf{X} \in \mathbb{S}^s,$$

\uparrow Linear SDP relaxation

Quadratic SDP: $\min c^T x$ sub to $M(x) \succeq \mathbf{O}$,

where $M : \mathbb{R}^s \rightarrow \mathbb{S}^n$ whose (i, j) element is given by

$$M_{ij}(x) = (1, x^T) Q_{ij} \begin{pmatrix} 1 \\ x \end{pmatrix} = Q_{ij} \bullet \begin{pmatrix} 1 & x^T \\ x & xx^T \end{pmatrix}, \forall x \in \mathbb{R}^s.$$

Here $Q \bullet Y = \text{trace } Q^T Y$ (the inner product of Q and Y).

(b) Linear SDP relaxation of a sparse quadratic SDP

SDP: $\min c^T x$ sub to $\widehat{M}(x, \mathbf{X}) \succeq \mathbf{O}, \begin{pmatrix} x_0 & x^T \\ x & \mathbf{X} \end{pmatrix} \succeq \mathbf{O}, x_0 = 1$

where $\widehat{M} : \mathbb{R}^s \times \mathbb{S}^s \rightarrow \mathbb{S}^n$ whose (i, j) element is given by


$$\widehat{M}_{ij}(x, \mathbf{X}) = Q_{ij} \bullet \begin{pmatrix} 1 & x^T \\ x & \mathbf{X} \end{pmatrix} \text{ for every } x \in \mathbb{R}^s, \mathbf{X} \in \mathbb{S}^s,$$

(b) Linear SDP relaxation of a sparse quadratic SDP

$$\text{SDP: } \min c^T x \text{ sub to } \widehat{M}(x, X) \succeq O, \begin{pmatrix} x_0 & x^T \\ x & X \end{pmatrix} \succeq O, x_0 = 1$$

where $\widehat{M} : \mathbb{R}^s \times \mathbb{S}^s \rightarrow \mathbb{S}^n$ whose (i, j) element is given by

$$\widehat{M}_{ij}(x, X) = Q_{ij} \bullet \begin{pmatrix} 1 & x^T \\ x & X \end{pmatrix} \text{ for every } x \in \mathbb{R}^s, X \in \mathbb{S}^s,$$



d-space sparsity ($\forall Q_{ij}$) and r-space sparsity (\widehat{M})
($s = 40, n = 41$)

(b) Linear SDP relaxation of a sparse quadratic SDP

$$\text{SDP: } \min c^T x \text{ sub to } \widehat{M}(x, X) \succeq O, \begin{pmatrix} x_0 & x^T \\ x & X \end{pmatrix} \succeq O, x_0 = 1$$

where $\widehat{M} : \mathbb{R}^s \times \mathbb{S}^s \rightarrow \mathbb{S}^n$ whose (i, j) element is given by

$$\widehat{M}_{ij}(x, X) = Q_{ij} \bullet \begin{pmatrix} 1 & x^T \\ x & X \end{pmatrix} \text{ for every } x \in \mathbb{R}^s, X \in \mathbb{S}^s,$$

(b) Linear SDP relaxation of a sparse quadratic SDP

$$\text{SDP: } \min c^T x \text{ sub to } \widehat{M}(x, X) \succeq O, \begin{pmatrix} x_0 & x^T \\ x & X \end{pmatrix} \succeq O, x_0 = 1$$

where $\widehat{M} : \mathbb{R}^s \times \mathbb{S}^s \rightarrow \mathbb{S}^n$ whose (i, j) element is given by

$$\widehat{M}_{ij}(x, X) = Q_{ij} \bullet \begin{pmatrix} 1 & x^T \\ x & X \end{pmatrix} \text{ for every } x \in \mathbb{R}^s, X \in \mathbb{S}^s,$$

		SDPA elapsed time in seconds			
s	n	no sparsity	d-space	r-space	d- & r-space
40	41	1.4	0.3	1.3	0.2
80	81	33.5	1.7	34.6	0.8
160	161	1427.1	19.6	1483.0	4.1
320	321	-	262.2	-	31.8

(c) SDP relaxation of POPs by SparsePOP+SDPA — 1
alkyl from globalib

$$\begin{aligned} \min & -6.3x_5x_8 + 5.04x_2 + 0.35x_3 + x_4 + 3.36x_6 \\ \text{sub.to} & -0.820x_2 + x_5 - 0.820x_6 = 0, \\ & 0.98x_4 - x_7(0.01x_5x_{10} + x_4) = 0, \\ & -x_2x_9 + 10x_3 + x_6 = 0, \\ & x_5x_{12} - x_2(1.12 + 0.132x_9 - 0.0067x_9^2) = 0, \\ & x_8x_{13} - 0.01x_9(1.098 - 0.038x_9) - 0.325x_7 = 0.574, \\ & x_{10}x_{14} + 22.2x_{11} = 35.82, \\ & x_1x_{11} - 3x_8 = -1.33, \text{ lbd}_i \leq x_i \leq \text{ubd}_i \ (i = 1, 2, \dots, 14). \end{aligned}$$

no sparsity		d-space sparsity	
E. time	ϵ_{obj}	E. time	ϵ_{feas}
> 10,000		1.3	8.2e-6 8.5e-10

ϵ_{obj} = approx. min. val. - lower bd. for the min. val.,

ϵ_{feas} = the max. error in equalities.

(c) SDP relaxation of POPs by SparsePOP+SDPA — 2
 Minimize the Broyden tridiagonal function $f_B(\mathbf{x})$ over \mathbb{R}^n .

$$f_B(\mathbf{x}) = \sum_{i=1}^n ((3 - 2x_i)x_i - x_{i-1} - 2x_{i+1} + 1)^2,$$

where $x_0 = 0$ and $x_{n+1} = 0$.

n	no sparsity	d-space	
	E. time	E. time	ϵ_{obj}
10	1.80	0.04	4.4e-9
20	916.95	0.08	1.5e-9
5000	o.mem.	29.44	5.1e-5
10000	o.mem.	59.52	9.2e-4

ϵ_{obj} = an approx. min. val. - a l. bound for the min. val..

Outline

- 0 Semidefinite Programming (SDP)
- 1 A simple example for 2 types of sparsities
- 2 Chordal graph
- 3 Domain-space sparsity
- 4 Range-space sparsity
- 5 Numerical results
- 6 Concluding remarks

Two types of sparsities of large-scale SDPs which are characterized by a chordal graph structure:

- (a) Domain-space sparsity
- (b) Range-space sparsity

- Numerical methods for converting large-scale SDPs into smaller SDPs by exploiting (a) and (b).

Linear,	each large-scale matrix variable
polynomial or	↓ exploiting (a) Domain-space sparsity
nonlinear	multiple smaller matrix variables
SDP	each large-scale matrix inequality
	↓ exploiting (b) Range-space sparsity
	multiple smaller matrix inequalities

Two types of sparsities of large-scale SDPs which are characterized by a chordal graph structure:

- (a) Domain-space sparsity
- (b) Range-space sparsity

- Numerical methods for converting large-scale SDPs into smaller SDPs by exploiting (a) and (b).

Linear,	each large-scale matrix variable
polynomial or	↓ exploiting (a) Domain-space sparsity
nonlinear	multiple smaller matrix variables
SDP	each large-scale matrix inequality
	↓ exploiting (b) Range-space sparsity
	multiple smaller matrix inequalities

- Very effective when SDP is sparse.
- Overheads in domain- & range-space conversion methods; adding equalities, real variables and/or matrix variables. Hence, less effective if SDP is denser.