

# Elementary Numerical Partial Differential Equations

## Homework 3

(Due: Jan. 24, 2008)

Consider the hyperbolic partial differential equation

$$\begin{aligned}u_t + cu_x &= 0, & -1 < x < 1, & \quad 0 < t < .5, \\u(1, t) &= u(-1, t), & 0 < t < .5, \\u(x, 0) &= v(x), & -1 < x < 1,\end{aligned}\tag{1}$$

where  $c > 0$ . Recall that the exact solution to (1) is given by

$$u(x, t) = v(x - ct)$$

Consider uniform refinement, that is, for  $h = 1/N$  and  $k = 1/M$ , we let  $x_j = jh$  and  $t_n = nk$ . Write a Matlab program to solve the equation with the Lax-Friedrichs method (13.5), Lax-Wendroff (13.17) method and Upwind method (13.22, 13.23).

- Consider the problem (1) with  $c = 0.5$ ,  $b(t) = 0$ , and

$$v(x) = \begin{cases} (x - 0.5)^2(x + 0.5)^2 \cdot 2^4 & -0.5 \leq x \leq 0.5 \\ 0 & \textit{otherwise} \end{cases}$$

Study the performance (stability and accuracy) of the schemes, compare the computed results with the exact solution and discuss the difference.