

• **Limits and Continuity:**

1. Let $f(x) = \begin{cases} |x+2| & \text{for } x \leq 0; \\ 2+x^2 & \text{for } 0 < x < 2; \\ x^3 & \text{for } x \geq 2 \end{cases}$.

Find (a) $\lim_{x \rightarrow 0^-} f(x)$, (b) $\lim_{x \rightarrow 0^+} f(x)$, (c) $\lim_{x \rightarrow 2^-} f(x)$, (d) $\lim_{x \rightarrow 2^+} f(x)$, (e) $\lim_{x \rightarrow 0} f(x)$, (f) $\lim_{x \rightarrow 2} f(x)$.

2. Compute $\lim_{x \rightarrow 0} \frac{1 - \cos 4x}{9x^2}$

3. Let $f(x) = \begin{cases} cx - 2 & \text{for } x \leq 2; \\ cx^2 + 2 & \text{for } x > 2 \end{cases}$ Find c such that $f(x)$ is continuous.

4. Determine the intervals on which $f(x) = \ln(1 - x^2)$ is continuous.

5. Compute

(i) $\lim_{x \rightarrow 0} \frac{\sqrt{x+9} - 3}{x}$

(ii) $\lim_{x \rightarrow 1^-} \frac{2x}{x^2 - 1}$

(iii) $\lim_{x \rightarrow \infty} \frac{x + \sin x}{4x + 999}$

• **Differentiation:**

1. Find the tangent line to the curve $y = x^3 - 4x^2 + 2x + 1$ at the point $(1, 0)$.

2. Let $y = e^{x^2} \sin(x^2 + x + 1) \cdot \sqrt{3x+1}/(x^2 - 1)$. Find $\frac{dy}{dx}$.

3. The equation $7x^2y^3 - 5xy^2 - 4y = 7$ defines y implicitly as a function of x . Find $\frac{dy}{dx}$.

4. Find the derivative of (i) $f(x) = x^{2x}$; (ii) $g(x) = \frac{x^2-x}{3x+1}$; (iii) $h(x) = \ln \sqrt{\frac{3x+1}{5x+2}}$ (assuming $3x+1 > 0$)

5. Compute $\frac{d}{dx} \cos^{-1}(2x^3)$

6. Determine if $f(x) = x^7 + 2x^3 - 2006$ is increasing, decreasing or neither. Prove $f(x) = 0$ has exactly one solution.

• **Application of Differentiation:**

1. Estimate $\tan((\pi/4) + 0.05)$ by the method of linear approximation (i.e., by differentials).

2. Compute $\lim_{x \rightarrow 1^+} \frac{\ln x}{(x-1)^2}$

3. Find the asymptotes of

(i) $f(x) = \frac{(3x-1)^2}{9x^2-4}$. (ii) $f(x) = \frac{(3x-1)^2}{9x^2-1}$. (iii) $f(x) = \frac{(3x-1)^2}{x-1}$

4. Let $f(x) = 2x^3 - 3x^2 - 12x$. Find the relative extrema of $f(x)$.

- Find the absolute maximum and minimum values of the function $f(x) = 2x^3 - 9x^2 + 12x$ over the interval $[0, 2]$.
- Determine the concavity of $f(x) = 4x^3 - x^4$.
- If 300 cm^2 of material is available to make a box with square base and an open top, find the largest possible volume of the box. Explain why your answer is the absolute maximum.
- Sketch the graph of the continuous function f that satisfies the conditions:

$$\begin{aligned} f''(x) &> 0 \quad \text{if } |x| > 2, & f''(x) < 0 & \quad \text{if } |x| < 2; \\ f'(0) &= 0, & f'(x) > 0, & \quad \text{if } x < 0, & f'(x) < 0, & \quad \text{if } x > 0; \\ f(0) &= 1, & f(2) = \frac{1}{2}, & & f(x) > 0 & \quad \text{for all } x, \text{ and } f \text{ is an even function.} \end{aligned}$$

- An automobile dealer is selling cars at a price of \$12,000. The demand function is $D(p) = 2(15 - 0.001p)^2$, where p is the price of a car. Should the dealer raise or lower the price to increase the revenue?
- Compute:
 - $\lim_{x \rightarrow 0} \left(\frac{1}{\ln(x+1)} - \frac{1}{x} \right)$
 - $\lim_{x \rightarrow 0^+} (\cos x)^{1/x}$
 - $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x} \right)^x$

• **Integration:**

- Let $f(x) = x + 1$
 - Divide the interval $[0, 5]$ into n equal parts, and using right endpoints find an expression for the Riemann sum R_n .
 - Using the answer you got from part(a), calculate $\lim_{n \rightarrow \infty} R_n$ (without using antiderivatives).

- Find the derivatives of the following functions. It is not necessary to simplify your answer: $G(x) = \int_0^{x^2} \sqrt{1+t^4} dt$

- Let f be continuous and define F by

$$F(x) = \int_0^x [t^2 \int_1^t f(u) du] dt.$$

Find $F'(x)$ and $F''(x)$.

- Evaluate the given integral

$$\begin{aligned} \text{(i)} & \int x(x+1)^9 dx, \quad \text{(ii)} \int \frac{\cos \theta}{\sin^2 \theta - 2 \sin \theta - 8} d\theta. \quad \text{(iii)} \int \frac{dx}{e^x \sqrt{4 + e^{2x}}}. \\ \text{(iv)} & \int \frac{\ln x}{x \sqrt{1 + \ln x}} dx, \quad \text{(v)} \int \frac{x^3}{\sqrt{x^2 + 1}} dx. \quad \text{(vi)} \int_{-\infty}^{\infty} \frac{e^{-x}}{(1 + e^{-x})^2} dx \end{aligned}$$

5. Evaluate the given integral
 (i) $\int \sqrt{x}e^{\sqrt{x}} dx$; (ii) $\int \frac{\sqrt{\ln x}}{x} dx$; (iii) $\int \frac{x}{x+4} dx$; (iv) $\int (\ln x)^2 dx$;
6. Evaluate the given integral
 (i) $\int \frac{\ln x}{x} dx$; (ii) $\int \ln(x^2) dx$; (iii) $\int \frac{3x}{x^2 - 3x - 4} dx$; (iv) $\int \frac{-2x^2 + 4}{x^3 + 2x^2 + x} dx$
7. Evaluate the definite integrals:
 (i) $\int_1^4 \sqrt{x}e^{\sqrt{x}} dx$; (ii) $\int_1^e (\ln x)^2 dx$;
8. Evaluate the given integral
 (i) $\int \sec^3 t dt$
 (ii) $\int \sec t dt$
9. (a) $\int_{-\infty}^{\infty} x^3 dx$ (b) $\lim_{R \rightarrow \infty} \int_{-R}^R x^3 dx$
10. Determine whether the integral converges or diverges:
 (i) $\int_0^1 x^{-1/3} dx$
 (ii) $\int_0^1 x^{-4/3} dx$
 (iii) $\int_1^{\infty} x^{-1/3} dx$
 (iv) $\int_{-1}^1 x^{-1/3} dx$

• **Application of Integration**

- Find the region bounded by the parabola $x = 2 - y^2$ and the line $y = x$.
- A solid is formed by revolving the circular disk $(x - 5)^2 + y^2 = 4$ about the y -axis. Set up, **but do not evaluate**, a definite integral which give the volume of the solid.
- Let Ω be the region bounded by $y = \sec x$, $x = 0$, $x = \frac{\pi}{4}$ and $y = 0$. Find integrals represent the volume of the solids generated by Ω about (a) x -axis, (b) y -axis, (c) $y = -1$, (d) $x = -1$.
(Don't evaluate the integrals)
- Set up a definite integral for the arc length of an ellipse $x^2 + 4y^2 = 4$.
- Set up the integral for the surface area of the surface of revolution. $y = e^x, 0 \leq x \leq 1$, revolved about x -axis.
- (i) At time t , a particle has position $x(t) = 1 - \cos t$, $y(t) = t - \sin t$ Find the total distance traveled from $t = 0$ to $t = 2\pi$. Find the speed of the particle at $t = \pi$.
 (ii) Find the area of the surface generated by revolving the curve $y = \cosh x = \frac{e^x + e^{-x}}{2}$, $x \in [0, \ln 2]$ about the x -axis.